

Deterministic Entanglement Swapping with an Ion Trap Quantum Computer

M. Riebe,¹ T. Monz,¹ K. Kim,^{1,*} A. S. Villar,^{2,†} P. Schindler,¹ M. Chwalla,¹ M. Hennrich,¹ and R. Blatt^{1,2}

¹*Institut für Experimentalphysik, Universität Innsbruck, Technikerstr. 25, A-6020 Innsbruck, Austria*

²*Institut für Quantenoptik und Quanteninformation, Österreichische Akademie der Wissenschaften, Technikerstr. 21A, A-6020 Innsbruck, Austria*

Entanglement – once only a subject of disputes about the foundation of quantum mechanics – has today become an essential issue in the newly emerged field of quantum information processing, promising such fascinating applications as secure communication, teleportation and powerful quantum computation. Accordingly, a focus of current experimental work in the quantum information field is the creation and manipulation of entangled quantum systems. Here, we present our results on entangling two qubits in an ion-trap quantum processor not through a direct interaction of the ion qubits but instead through the action of a protocol known as *entanglement swapping*[1]. Our ion-trap system allows us to implement all steps of the entanglement swapping protocol in a fully deterministic way. Thus, two ion-qubits can be prepared on demand in a well-defined entangled state. This particular feature may facilitate the implementation of quantum repeaters[2] or aid in distributing entangled states in ion-trap quantum computers[3].

Entanglement among quantum systems usually results from a common origin of the systems or a direct local interaction between the systems. In both cases the entangling operation requires the systems to be at the same point in space at a certain time. In entanglement swapping, however, the entangling operation is achieved as follows: Starting from two entangled pairs, A_1A_2 and B_1B_2 , a joint measurement of A_2 and B_2 in the Bell basis projects pair A_1B_1 in an entangled state even though A_1 and B_1 might be far apart. This can be used for applications such as quantum communication or teleportation, which require that quantum systems at distant locations are entangled. Entanglement swapping facilitates the provision of such separated pairs of entangled systems, as two quantum systems in two distinct locations can be entangled via a chain of intermediary entangled pairs. Furthermore, if entanglement swapping is combined with entanglement purification, as proposed in quantum repeater schemes[2], the final entangled state can have a higher fidelity than is achievable with a direct transfer of an entangled system. Up to now, entanglement swapping has been implemented with photon pairs[4, 5] and with ion-photon entangled states[6]. These implementations are appealing as the photons can be easily used to transfer quantum information over large distances. However, in these experiments successful entanglement swapping events could only be established probabilistically for a small subset of all states. By contrast, quantum information processing with a system of trapped ions has the advan-

tage that all basic operations for handling quantum information are implemented with high fidelities and, in particular, in a completely deterministic fashion. This capability has been demonstrated in various experiments, for example by implementing a completely deterministic quantum teleportation protocol between trapped-ion qubits[7, 8]. Here we apply the techniques of these experiments to implement an entanglement swapping protocol for four trapped ion qubits. In contrast to a previous realization with trapped ions which generated an entangled state at a distance of approximately one meter[6] this realization is fully deterministic. Entanglement purification, a protocol equally important for the generation of distant entanglement, has been demonstrated in[14]. This experiments shows a similar level of control of four ions as our experiment, but does not feature a deterministic control of the final state.

In the experiments described here, deterministic entanglement swapping is implemented with strings of $^{40}\text{Ca}^+$ ions confined in a linear Paul trap. Each ion represents a qubit, where quantum information is stored in a superposition of the $|S\rangle \equiv S_{1/2}(m = -1/2)$ ground state and the meta-stable $|D\rangle \equiv D_{5/2}(m = -1/2)$ state. Additionally, the ion string undergoes a harmonic motion in the trap potential. For our purpose, we only consider the centre-of-mass (COM) vibrational mode along the trap axis at a frequency of $\omega_z = 2\pi \cdot 1.2$ MHz. Laser cooling and optical pumping allow us to prepare the COM mode in its ground state $|n = 0\rangle$ and to initialize all qubits in the state $|S\rangle$ [9]. After initialization to $|SS\dots S\rangle \otimes |0\rangle \equiv |SS\dots S, 0\rangle$, the quantum state of the ion string is manipulated by applying laser pulses resonant with the $S_{1/2} \leftrightarrow D_{5/2}$ transition to individual ions. This interaction is precisely controlled by setting the frequency, length, phase and intensity of the pulses. Two kinds of operations are used: a) Laser pulses at the resonance frequency, ω_0 , of the $S_{1/2} \leftrightarrow D_{5/2}$ transition which implement single qubit rotations and b) laser pulses at the blue sideband frequency, $\omega_0 + \omega_z$, which change the electronic state of the addressed ion as well as the vibrational state of the ion string. The latter kind of operation enables us to implement quantum gates between arbitrary ions within the ion string mediated by the vibrational mode. The quantum state of the qubits is measured by scattering light on the $S_{1/2} \leftrightarrow P_{1/2}$ transition at 397 nm and observing the resulting resonance fluorescence with a CCD camera. This corresponds to a projective measurement of the qubits in the $\{|S\rangle, |D\rangle\}$ basis, where the observation of resonance fluorescence at 397 nm indicates a projection to the $|S\rangle$ -state, while the absence of fluorescence indicates a projection to the $|D\rangle$ -state. Further

details of the experimental setup and methods are given in Ref. [9].

Entanglement swapping is realized in our experimental setup as shown in Fig. 1. Initially a string of four $^{40}\text{Ca}^+$ ions is prepared in the state $|SSSS, 0\rangle$ (the order of the ions is |ion 4...ion 2 ion 1, 0>). First, two pairs of entangled states are prepared. This is achieved by a sequence of three laser pulses which is first applied to ions 1 and 2, and later to ions 3 and 4, and which prepares each pair in the state $|\Psi_{ij}^-\rangle = (|DS\rangle_{ij} - |SD\rangle_{ij})/\sqrt{2}$ with $i, j \in \{1, \dots, 4\}$. The total state of the ion string $|\Psi_{1234}\rangle = |\Psi_{12}^-\rangle \otimes |\Psi_{34}^-\rangle$ is rewritten by grouping the unentangled ions 2,3 and 1,4 into pairs and expressing their state in terms of the Bell states $|\Phi^\pm\rangle = (|DD\rangle \pm |SS\rangle)/\sqrt{2}$ and $|\Psi^\pm\rangle = (|DS\rangle \pm |SD\rangle)/\sqrt{2}$ as:

$$|\Psi_{1234}\rangle = -\frac{1}{2} \left(|\Phi_{14}^+\rangle \otimes |\Phi_{23}^+\rangle - |\Phi_{14}^-\rangle \otimes |\Phi_{23}^-\rangle - |\Psi_{14}^+\rangle \otimes |\Psi_{23}^+\rangle + |\Psi_{14}^-\rangle \otimes |\Psi_{23}^-\rangle \right). \quad (1)$$

Writing the state in this way, the procedure for entanglement swapping can be directly inferred: After a projective measurement of either pair (1,4 or 2,3) in the Bell basis the other pair will be in an entangled state; thus two ions become entangled which have not previously interacted. The required Bell measurement is achieved in our experiment by mapping the Bell state basis, $\{|\Phi_\pm\rangle, |\Psi_\pm\rangle\}$, onto the product state basis, $\{|DD\rangle, |DS\rangle, |SD\rangle, |SS\rangle\}$, in which a projective measurement is easily possible in our experimental setup. Consider for example that the Bell measurement is applied to ions 1,4 in order to entangle ions 2,3 (see Methods). This map is implemented by a controlled-NOT (CNOT) gate [10], with ion 1 as the control and ion 4 as the target qubit, followed by a $\pi/2$ -rotation on the control qubit. After this step the total state of the ion string is given by:

$$|\Psi_{1234}\rangle = \frac{1}{2} \left(|DD\rangle_{14} \otimes |\Phi_{23}^+\rangle + e^{i\phi} |DS\rangle_{14} \otimes |\Phi_{23}^-\rangle - e^{-i\phi} |SD\rangle_{14} \otimes |\Psi_{23}^+\rangle - |SS\rangle_{14} \otimes |\Psi_{23}^-\rangle \right) \quad (2)$$

The additional phase factors, $\exp(i\phi)$ with $\phi = \pi/\sqrt{2}$, are caused by the CNOT gate but are ignored due to the subsequent projective measurement. The states of ions 1 and 4 are measured by exposing the ion string for 300 μs with light at 397 nm and observing the resonance fluorescence of the ions with a photomultiplier tube (PMT). The quantum information stored in the other ions is protected from the 397 nm light during the measurements (see Methods). From (2) it can be seen that the observed states of ions 1 and 4 exactly correlates with the entangled state of ions 2 and 3 (either Φ^+ , Φ^- , Ψ^+ , Ψ^-). This fact

allows us to create a well-defined entangled output state, which is a requirement for further use of this newly entangled state in protocols like teleportation or purification. The result of this Bell measurement determines which single qubit rotation is to be applied to ions 2 or 3 in order to obtain the desired entangled state. We choose $|\Psi_{23}^-\rangle$ as the desired output state and note that $|\Psi^-\rangle = I \otimes Z|\Psi^+\rangle = I \otimes X|\Phi^-\rangle = I \otimes XZ|\Phi^+\rangle$, where I is the identity operator and X and Z are Pauli-operators. To map ions 2,3 deterministically to $|\Psi^-\rangle$, first the fluorescence counts measured with the PMT are evaluated and the states of ions 1 and 4 are determined. Then depending on this result, laser pulses are applied to ion 3, which implement the required X - and Z -operations.

The protocol described above is experimentally implemented by the sequence of laser pulses given in Tab. (I). To prove that this sequence properly implements entanglement swapping, we analyze the output state ρ_{23} of ions 2 and 3 by quantum state tomography. This requires that a complete tomographic data set is recorded, from which ρ_{23} is reconstructed[11]. First, to check whether the protocol is working as expected, we omit the conditional rotations and measure the quantum state directly after the rotation of the Bell basis onto the product basis. At this point of the protocol the ion string is expected to be in the state given in (2). Here, the reconstruction of ρ_{23} would only yield a completely mixed state, if the result of the Bell measurement on ions 1 and 4 is neglected. However, sorting the tomographic data set with respect to the result of the Bell measurement and then reconstructing ρ_{23} for each data subset yields the result shown in Fig. (2). For each Bell measurement result we observe ions 2 and 3 to be in the entangled state expected from (2) with fidelities ranging between 76% and 91%, and an entanglement of formation between 0.49 and 0.75[12]. This is sufficient to prove that the entanglement swapping protocol is working, but requires a postprocessing procedure as described above. On the other hand, by implementing the complete protocol including the conditional rotations our scheme becomes completely deterministic, and the final state of ions 2 and 3 will always be $|\Psi^-\rangle$ irrespective of the Bell measurement result. Figure (3) shows the measured density matrix of ions 2 and 3 for this case. For the reconstruction of this density matrix the state of ions 1 and 4 was neglected and no further postprocessing of the data set was applied. Indeed, ions 2 and 3 are deterministically mapped onto state $|\Psi^-\rangle$ with a fidelity of $\langle\Psi^-|\rho_{23}|\Psi^-\rangle = 79(2)\%$. In addition, this state violates a Clauser-Horne-Shimony-Holt inequality [13] as $\langle A \rangle = 2.18(8) > 2$, where $A = (\sigma_x \otimes \sigma_{z-x} - \sigma_x \otimes \sigma_{z+x} - \sigma_z \otimes \sigma_{z+x} - \sigma_z \otimes \sigma_{z-x})/\sqrt{2}$ with $\sigma_{z\pm x} = \sigma_z \pm \sigma_x$.

The achieved fidelity is mainly limited by the imperfect fidelities of the initial Bell states ($\approx 96\%$), and of the CNOT gate operation ($\approx 93\%$ [10]).

In summary, we have demonstrated how ion qubits that never interacted are entangled through the action of entanglement swapping. In particular, we have realized this scheme in a completely deterministic manner. Within the bounds of the measured fidelity we achieve entanglement swapping in every single experimental run and then prepare the target ions in a well-defined entangled state. This scheme will find various applications in the future, for example in an implementation of quantum repeaters as proposed in [2]. For this, the presented entanglement swapping protocol would be combined with entanglement purification [14] to generate entangled qubit pairs of high fidelity at distant locations. A practical implementation of this scheme will require an efficient photon-ion qubit interface[15, 16]. Another application of the entanglement swapping protocol will be in large-scale ion-trap quantum computers [3], where it can help to prepare entangled pairs of qubits for quantum teleportation, to connect distant regions of the quantum processor, while minimizing the need to transport ions between distinct trap locations.

Methods

In order to achieve the reported entanglement swapping fidelity, we have to take several measures, to counteract possible error sources in our experimental setup. A major concern are addressing errors, i.e. the fact that every laser pulse not only manipulates the correct ion but also carries out an unwanted operation on the neighbouring ion(s)[9]. This error source motivates the choice of ions 1 and 4 for the Bell measurement, such that a major part of the pulse sequence is carried out on the outer ions, where addressing errors are smaller as only one neighbouring ion is present.

During the measurement of either ion 1 or 4, the quantum information in all other ions has to be preserved from the destructive influence of the 397 nm light. For this purpose we shift the S -state population of these ions to the Zeeman level $D' = D_{5/2}(m = -5/2)$, such that the quantum information is completely encoded in the $D_{5/2}$ -state[17]. Ions 2 and 3 are transferred to this state directly after the initial Bell states are prepared. This transfer also minimizes addressing errors due to laser pulses applied to ions 1 and 4. A problem arises because quantum information encoded in the D -state is highly susceptible to phase shifts caused by magnetic field variations. We counter this problem by applying a spin-echo sequence after approximately half of the pulse sequence to cancel out phase shifts due to low-frequency magnetic field changes[18].

-
- * Current Address: Department of Physics and Joint Quantum Institute, University of Maryland, College Park, Maryland, 20742, USA
- † Current Address: Institute of Optics, Information and Photonics, Max Planck Research Group, University of Erlangen-Nuernberg, Staudtstrasse 7/B2, 91058 Erlangen, Germany
- [1] Żukowski, M., Zeilinger, A., Horne, M. A. & Ekert, A. K. “Event-ready-detectors” Bell experiment via entanglement swapping. *Phys. Rev. Lett.*, **71**, 4287–4290 (1993).
- [2] Briegel, H.-J., Dür, W., Cirac, J. I. & Zoller, P. Quantum repeaters: The role of imperfect local operations in quantum communication. *Phys. Rev. Lett.*, **81**, 5932–5935 (1998).
- [3] Kielpinski, D., Monroe, C. & Wineland, D. J. Architecture for a large-scale ion-trap quantum computer. *Nature*, **417**(6890), 709–711 (2002).
- [4] Pan, J.-W., Bouwmeester, D., Weinfurter, H. & Zeilinger, A. Experimental entanglement swapping: Entangling photons that never interacted. *Phys. Rev. Lett.*, **80**, 3891–3894 (1998).
- [5] Halder, M. *et al.* Entangling independent photons by time measurement. *Nature Physics*, **3**(10), 692–695 (2007).
- [6] Moehring, D. L. *et al.* Entanglement of single-atom quantum bits at a distance. *Nature*, **449**(7158), 68–71 (2007).
- [7] Riebe, M. *et al.* Deterministic quantum teleportation with atoms. *Nature*, **429**(6993), 734–737 (2004).
- [8] Barrett, M. D. *et al.* Deterministic quantum teleportation of atomic qubits. *Nature*, **429**(6993), 737–739 (2004).
- [9] Schmidt-Kaler, F. *et al.* How to realize a universal quantum gate with trapped ions. *Appl. Phys. B*, **77**, 789–796 (2003).
- [10] Riebe, M. *et al.* Process tomography of ion trap quantum gates. *Phys. Rev. Lett.*, **97**, 220407 (2006).
- [11] Roos, C. F. *et al.* Bell states of atoms with ultralong lifetimes and their tomographic state analysis. *Phys. Rev. Lett.*, **92**(22), 220402 (2004).
- [12] Wootters, W. K. Entanglement of formation of an arbitrary state of two qubits. *Phys. Rev. Lett.*, **80**, 2245–2248 (1998).
- [13] Clauser, J. F., Horne, M. A., Shimony, A. & Holt, R. A. Proposed experiment to test local

- hidden-variable theories. *Phys. Rev. Lett.*, **23**, 880–884 (1969).
- [14] Reichle, R. *et al.* Experimental purification of two-atom entanglement. *Nature*, **443**(7113), 838–841 (2006).
- [15] Boozer, A. D., Boca, A., Miller, R., Northup, T. E. & Kimble, H. J. Reversible state transfer between light and a single trapped atom. *Physical Review Letters*, **98**(19), 193601 (2007).
- [16] Cirac, J. I., Zoller, P., Kimble, H. J. & Mabuchi, H. Quantum State Transfer and Entanglement Distribution among Distant Nodes in a Quantum Network. *Phys. Rev. Lett.*, **78**, 3221–3224 (1997).
- [17] Roos, C. F. *et al.* Control and measurement of three-qubit entangled states. *Science*, **304**(5676), 1478–1480 (2004).
- [18] Hahn, E. L. Spin echoes. *Phys. Rev.*, **80**, 580–594 (1950).

Addendum We gratefully acknowledge support by the Austrian Science Fund (FWF), by the European Commission (SCALA, CONQUEST networks), and by the Institut für Quanteninformation GmbH. This material is based upon work supported in part by the U. S. Army Research Office.

The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to Mark Riebe. (email: Mark.Riebe@uibk.ac.at).

Step	Pulse	Description
1	$R_1^+(\pi/2, \pi/2)$	Entangle ion 1 and 2
2	$R_2^C(\pi, -\pi/2)$	
3	$R_2^+(\pi, \pi/2)$	
4	$R_3^+(\pi/2, \pi/2)$	Entangle ion 3 and 4
5	$R_4^C(\pi, -\pi/2)$	
6	$R_4^+(\pi, \pi/2)$	
7	$R_2^H(\pi, 0)$	Hide ion 2 and 3
8	$R_3^H(\pi, 0)$	
9	$R_1^+(\pi, 0)$	CNOT gate
10	$R_4^C(\pi/2, -\pi/2)$	
11	$R_4^+(\pi/2, \pi/2)$	
12	$R_4^+(\sqrt{2}\pi, 0)$	
13	$R_4^+(\pi/2, -\pi/2)$	
14	$R_4^C(\pi/2, \phi)$	
15	$R_1^+(\pi, \pi)$	
16	$R_2^C(\pi, 0)$	Spin echo ion 2
17	$R_2^H(\pi, 0)$	
18	$R_2^C(\pi, 0)$	
19	$R_3^C(\pi, 0)$	Spin echo ion 3
20	$R_3^H(\pi, 0)$	
21	$R_3^C(\pi, 0)$	
22	$R_1^C(\pi/2, -\phi)$	Finish Bell rotation
23	$R_4^H(\pi, 0)$	Hide ion 4
24	397 nm light	Measure ion 1 (300 μ s)
25	$R_4^H(\pi, \pi)$	Undo hide ion 4
26	$R_1^H(\pi, 0)$	Hide ion 1
27	397 nm light	Measure ion 4 (300 μ s)
28	$R_1^H(\pi, \pi)$	Undo hide ion 1
29	$R_2^H(\pi, \pi)$	Undo hide ion 2
30	$R_3^H(\pi, \pi)$	Undo hide ion 3
31	$R_3^C(\pi, 0)$	Phase flip of ion 3 if ion 1 in $ D\rangle$
32	$R_3^C(\pi, \pi/2)$	
33	$R_3^C(\pi, 0)$	Bit flip of ion3 if ion 4 in $ D\rangle$

TABLE I: Sequence of laser pulses and experimental steps to implement entanglement swapping. Laser pulses applied to the i -th ion on carrier transitions are denoted by $R_i^C(\theta, \varphi)$ and $R_i^H(\theta, \varphi)$ and pulses on the blue sideband transition by $R_i^+(\theta, \varphi)$, where $\theta = \Omega t$ is the pulse area in terms of the Rabi frequency, Ω , the pulse length, t , and its phase, φ [11]. The index C denotes carrier transitions between the two logical eigenstates, while the index H labels transitions from the $S_{1/2^-}$ to the additional $D_{5/2^-}$ -sublevel used to hide individual ion qubits.




FIG. 1: Schematic showing the steps of the entanglement swapping protocol. Ion 2 and 3 are mapped to the state Ψ^- by a Bell measurement on ion 1,4 and subsequent conditional rotations. During the sequence the quantum information of some qubits is protected from 397 nm light by shifting their S-state population to the D-manifold (see Methods), denoted in the scheme as "Hide" and the inverse operation as "Hide⁻¹". The time during which the information is hidden is indicated by a grey line.




FIG. 2: Result of entanglement swapping without the final conditional rotations. The tomographic data set has been sorted with respect to the result of the Bell measurement. For each of the four subsets the density matrix of ions 2 and 3 was reconstructed. The real part of the obtained density matrices for the Bell measurement results $|DD\rangle$, $|DS\rangle$, $|SD\rangle$ and $|SS\rangle$ is shown in a), b), c) and d), respectively. The density matrices show ion 2 and 3 to be in the state expected from equation (2), with fidelities of a) 76(5)%, b) 82(5)%, c) 81(5)% and d) 91(3)%. Here, the error is due to statistical uncertainties in the tomographic measurements [11].




FIG. 3: Result of the fully deterministic entanglement swapping protocol. State Ψ^- was chosen as the desired output state. Here only the real part of the reconstructed density matrix ρ_{23} is shown. The experimentally achieved overlap with the ideal output state Ψ^- is $F = \langle \Psi^- | \rho_{23} | \Psi^- \rangle = 79(3)\%$.